DATOS SOBRE

EL EMBARAZO Y LA VIOLENCIA DOMÉSTICA

PORQUE ES IMPORTANTE

La violencia ocurre comúnmente durante el embarazo: entre el 4 y el 8 por ciento de las mujeres sufren violencia doméstica durante el embarazo.[1] Los efectos de la violencia durante el embarazo pueden ser devastadores tanto para la madre como para el feto. La violencia doméstica durante el embarazo está relacionado con la depresión, el abuso de sustancias, el uso de tabaco, la anemia, el sangrado del primer y segundo trimestre, y una reducción en el peso al nacer.[2] Desafortunadamente, este riesgo mortal para la salud a menudo no se detecta durante el embarazo debido a que la violencia doméstica rara vez se examina durante los exámenes prenatales.

¿SABÍAS QUE?

- Cada año, aproximadamente 1.5 millones de mujeres en los Estados Unidos son violadas o agredidas físicamente por su pareja.[3] Este número incluye a más de 324,000 mujeres que estaban embarazadas cuando ocurrió la violencia.[4]
- Entre las mujeres cuyos embarazos fueron planeados, el 5.3% reportó abuso durante el embarazo, en comparación con el 12.6% de las mujeres cuyos embarazos no fueron planeados y 15.3% para mujeres cuyos embarazos no fueron deseados.[5]
- 50-70% de las mujeres abusadas antes del embarazo, son abusadas durante el embarazo.
- El 26% de las adolescentes embarazadas informaron haber sido abusadas físicamente por sus novios. Casi la mitad de ellos dijeron que los golpes comenzaron o se intensificaron después de enterarse sobre el embarazo.[7]

HOMICIDIO Y EMBARAZO

- El asesinato es la segunda causa más común de muerte en mujeres embarazadas después de accidentes automovilísticos relacionada con lesiones durante el embarazo (31%).[8]
- Entre 1990 y 2004, más de 1,300 mujeres embarazadas fueron asesinadas.[9] La mayoría de estas mujeres (56%) fueron asesinadas a tiros, mientras que el resto fueron apuñaladas o estranguladas.[10]
- El 77% de las víctimas de homicidio embarazadas mueren durante el primer trimestre del embarazo.[11]
- Muchas muertes asociadas con el embarazo pueden pasar desapercibidas porque los certificados de muerte y los registros de médicos forenses no siempre indican si la fallecida estaba embarazada.[12]

QUIÉN ESTÁ MÁS EN RIESGO

- Las mujeres menores de 20 años y las mujeres que reciben atención prenatal tardía o no reciben atención prenatal son más vulnerables a homicidio por parte de su pareja.[3]
- Las mujeres con embarazos no deseados tienen de dos a cuatro veces más probabilidades de sufrir violencia física que las mujeres con embarazos planificados.[14]

EL IMPACTO DE LA VIOLENCIA DOMÉSTICA EN EL EMBARAZO

- Las mujeres que sufren abusos durante el embarazo tienen más probabilidades de retrasar el ingreso a la atención prenatal.[15]
- Complicaciones del embarazo, incluyendo poco aumento de peso, anemia, infecciones y primer y segundo trimestre sangrado, son significativamente mayores para las víctimas de violencia doméstica.[16]
- Las mujeres embarazadas que son víctimas de la violencia de su pareja íntima tienen más probabilidad de sufrir depresión y suicidio, así como consumir tabaco, alcohol y drogas durante el embarazo.[17]

EXAMEN DE ATENCIÓN MÉDICA

- El 96% de las mujeres en los Estados Unidos que dan a luz a un bebe vivo reciben atención prenatal.[18] Las mujeres embarazadas tienen un promedio de 12 a 13 visitas durante el embarazo.[19] A pesar de esta interacción frecuente, menos de la mitad de los proveedores de salud hablan con sus pacientes sobre violencia de pareja.[20]
- Solo el 18% de las mujeres embarazadas examinadas en la sala de emergencias reportan que el médico les preguntó sobre violencia infligida por la pareja.[21]

SI NECESITAS AYUDA

Si usted o alguien que conoce es víctima de violencia doméstica y desea ayuda, llame a la Linea de Ayuda Nacional de Violencia Doméstica al 1-800-799-SAFE.

Obtenga la ayuda que usted y su bebe necesitan para tener un comienzo sano. Llame a Healthy Start al 210-207-4725.

Te queremos. Queremos que estés aquí.
sanantonio.gov/violenceprevention

LOVE IS...

CITY OF SAN ANTONIO
METROPOLITAN HEALTH DISTRICT
VIOLENCE PREVENTION

FUENTES

[1]Gazamarian, JA, et al., "Violence and Reproductive Health: Current Knowledge and Future Research Directions", Maternal and Child Health Journal, Vol. 4, No 2, pg. 80 (2000). [2]Parker, B., McFarlane, J., & Soeken, K., "Abuse During Pregnancy: Association with Maternal Health and Infant Birthweight," Nursing Research, No. 45, pg. 32-37 (1996); McFarlane, J. et. Al. "Physical Abuse, Smoking and Substance Abuse During Pregnancy: Prevalence, Interrelationships and Effects on Birthweight," Journal of Obstetrical Gynecological and Neonatal Nursing, Vol. 25, pg. 313-20 (1996). [3]U.S. Department of Justice, Extent, Nature, and Consequences of Intimate Partner Violence: Research Report iii (2000). [4]Gazamarian, JA, et al., "Violence and Reproductive Health: Current Knowledge and Future Research Directions", Maternal and Child Health Journal, Vol. 4, No 2, pg. 80 (2000). [5]Parsons, L., Goodwin, M.M., and Petersen, R. "Violence Against Women and Reproductive Health: Toward Defining a Role for Reproductive Health Care Services," Maternal and Child Health Journal, Vol. 4, No. 2. pg. 135 (2000). [6]Saltzman, LE., Johnson, C.H., Gilbert, BC., and Goodwin, F. "Physical Abuse Around the Time of Pregnancy: An Examination of Prevalence and Risk Factors in 16 States. Maternal and Child Health Journal, Vol. 7, pg. 31-42 (2003). [7]Brustin, S., "Legal Response to Teen Dating Violence," Family Law Quarterly, vol. 29, no. 2, 333-334 (Summer 1995) (citing Worcester, "A More Hidden Crime: Adolescent Battered Women," The Network News, July/Aug., national Women's Health Network 1993). [8]Jeani Chang, MPH, Cynthia J. Berg, MD, MPH, Linda E. Saltzman, PhD and Joy Herndon, MS, "Homicide: A Leading Cause of Injury Deaths Among Pregnant and Postpartum Women in the United States, 1991-1999," American Journal of Public Health, Vol. 95, No. 3, pg. 471-77 (2005). [9]St. George, Donna, "Pregnant Women Murdered at an Alarming Rate," The Washington Post, Dec. 19, 2004. [10]Id. [11]Krulewitch, C., Pierre-Louis, M.L., deLeon-Gome, R., Guy, R., and Green, R. "Hidden From View: Violent Deaths Among Pregnant Women in the District of Columbia, 1988-1996". Journal of Midwifery and Women's Heath. Vo.l 46, pg 7 (2001). [12]Horon, I. & Cheng D., "Enhanced Surveillance for Pregnancy-Associated Mortality-Maryland, 1993-1998," The Journal of the American Medical Association, 285, No. 11, March 21, 2001. [13]Jeani Chang, MPH, Cynthia J. Berg, MD, MPH, Linda E. Saltzman, PhD and Joy Herndon, MS, "Homicide: A Leading Cause of Injury Deaths Among Pregnant and Postpartum Women in the United States, 1991-1999," American Journal of Public Health, Vol. 95, No. 3, pg. 471-77 (2005). [14]Gazamarian, JA, et al., "Violence and Reproductive Health: Current Knowledge and Future Research Directions", Maternal and Child Health Journal, Vol. 4, No 2, pg. 80 (2000). [15]Parker, B., "Physical and Emotional Abuse in Pregnancy: A Comparison of Adult & Teenage Women,"Nursing Research, Vol 4, No.2, pg. 80 (2000). [16]Parker, B., McFarlane, J., & Soeken, K., "Abuse During Pregnancy: Association with Maternal Health and Infant Birthweight," Nursing Research, No. 45, pg. 32-37 (1996). [17]McFarlane, J. et. Al. "Physical Abuse, Smoking and Substance Abuse During Pregnancy: Prevalence, Interrelationships and Effects on Birthweight," Journal of Obstetrical Gynecological and Neonatal Nursing, Vol. 25, pg. 313-20 (1996). [18]Ventura SJ, Martin JA, Curtin SC, Mathews TJ., "Report of Final Natality Statistics," Monthly Vital Statistics Report; Vol. 46, No.11 (suppl), National Center for Health Statistics (1998). [19]Id. [20]Parsons, L., Goodwin, M.M., and Petersen, R. "Violence Against Women and Reproductive Health: Toward Defining a Role for Reproductive Health Care Services," Maternal and Child Health Journal, Vol. 4, No. 2. pg. 135 (2000). [21]McGrath, ME., Hogan, JW., Peipert, JF. "A Prevalence Survey of Abuse and Screening for Abuse in Urgent Care Patients. Obstetric Gynecology,Vol. 91(4) pg 511-4. 1991.